

## Erratum

# Functional and molecular characterisation of mammary side population cells

Azra J Alvi<sup>1</sup>, Helen Clayton<sup>1</sup>, Chirag Joshi<sup>2</sup>, Tariq Enver<sup>2</sup>, Alan Ashworth<sup>1</sup>, Maria d M Vivanco<sup>1</sup>, Trevor C Dale<sup>3</sup>, Matthew J Smalley<sup>3,4</sup>

<sup>1</sup>Breakthrough Toby Robins Breast Cancer Centre, Institute of Cancer Research, London, UK

<sup>2</sup>Section of Gene Function and Regulation, Institute of Cancer Research, London, UK

<sup>3</sup>Section of Cell and Molecular Biology, Institute of Cancer Research, London, UK

<sup>4</sup>Current address: Division of Molecular Genetics, Netherlands Cancer Institute, Amsterdam, The Netherlands

Published: 4 December 2002

*Breast Cancer Res* 2003, **5**:E1 (DOI 10.1186/bcr563)

© 2003 BioMed Central Ltd (Print ISSN 1465-5411; Online ISSN 1465-542X)

## Erratum

It has been brought to our attention that there was an inaccuracy in the above article [1]. In the discussion section the authors state that “One of the assumptions underlying mammary transplant studies is that mammary outgrowths at limiting dilution are clonal (14,000–20,000 cells per cleared fat pad)” and they attribute this to reference 21 “Smith GH: Experimental mammary epithelial morphogenesis in an in vivo model: evidence for distinct cellular progenitors of the ductal and lobular phenotype. *Breast Cancer Res Treat* 1996, **39**:21-31” [2]. In the paper by Smith no claim of clonality is made, rather it is stated that it is not known whether the simultaneous development of both lobular and ductal phenotypes indicates cooperative interaction between the two epithelial progenitors or signals the presence of a third progenitor type capable of producing both ductular and lobular committed daughters. It is, therefore, incorrect that the assumption of clonality based on limiting dilution has been attributed to Gilbert Smith.

## References

1. Alvi A, Clayton H, Joshi C, Enver T, Ashworth A, Vivanco MdM, Dale TC, Smalley MJ: **Functional and molecular characterisation of mammary side population cells.** *Breast Cancer Res* 2003, **5**:R1-R8
2. Smith GH: **Experimental mammary epithelial morphogenesis in an in vivo model: evidence for distinct cellular progenitors of the ductal and lobular phenotype.** *Breast Cancer Res Treat* 1996, **39**:21-31