

VIEWPOINT

# Slugging their way to immortality: driving mammary epithelial cells into a stem cell-like state

Kelly Soady<sup>1</sup> and Matthew J Smalley<sup>\*2</sup>

## Abstract

Delineating the molecular factors that define and maintain the mammary stem cell state is vital for understanding normal development and tumorigenesis. A recent study by Guo and colleagues identifies two master transcriptional regulators of mammary stem cells, *Slug* and *Sox9*, ectopic expression of which confers stem cell attributes on differentiated mammary epithelial cells. *Slug* and *Sox9* expression was also shown to determine *in vivo* metastatic potential of human breast cancer cell lines. Understanding these factors in the context of normal lineage differentiation is an important step toward elucidating the mammary epithelial cell hierarchy and the origins of cancer stem cells.

## Background

The mammary gland is a highly dynamic tissue, undergoing significant morphological change during puberty, pregnancy, lactation and involution. Flow cytometry using multiple markers has enabled the prospective isolation of mammary epithelial subpopulations suggesting a hierarchical organisation of stem, progenitor and differentiated cells. Utilising the cleared mammary fat pad transplantation assay, developed from the pioneering work of Deome and colleagues [1], mammary stem cells (MaSCs) have been defined as highly enriched in the  $CD24^{+}/LowSca1-CD49^{High}CD29^{High}$  population [2-5]. A single cell from this population is able to recapitulate the entire gland upon transplantation [3,5], showing defining stem cell characteristics of multi-differentiative potential and self-renewal. Despite much work to define factors necessary for MaSC function, molecular mechanisms regulating MaSCs are poorly

defined. Epithelial–mesenchymal transition (EMT), a key developmental programme in the embryo that confers mesenchymal cell traits on epithelial cells, has recently been linked to the MaSC state [6,7]. EMT has also been implicated in tumour invasion and metastasis, and a potential role in self-renewal strengthens known associations between normal tissue stem cells and cancer cells [8].

Despite ongoing controversy over the existence and origin of cancer cells, the cancer stem cell model – in which tumours are maintained by a population of stem-like cancer cells – provides an attractive framework to understand metastatic potential and tumour heterogeneity in response to treatment. Irrespective of the origins of these cells, their exact nature and nomenclature and their operational detection, it is unlikely that neoplastic (stem-like) cell populations invent a novel programme to drive their sustained proliferation. Instead, cancer cells adopt the self-renewal programme active in the antecedent stem cell population and exploit this to organise the complex tissues observed at various stages of neoplastic progression. Delineating the molecular factors that define the mammary epithelial cell hierarchy and maintain the MaSC state is therefore an important step towards understanding both normal development and tumorigenesis.

## Slug and Sox9 as master regulators of the mammary stem cell state

A recent study by Guo and colleagues identified two master transcriptional regulators of the mammary stem cell state: the EMT-associated transcription factor *Slug*, and the SRY-box transcription factor *Sox9* [9]. Ectopic co-expression of *Slug* and *Sox9* in primary mouse mammary cells *in vitro* for 5 days was sufficient to convert differentiated luminal cells into MaSCs, as demonstrated by a competitive cleared mammary fat pad transplantation assay. Differentiated luminal cells expressing *Slug* and *Sox9* were able to reconstitute the entire mammary gland upon serial transplantation, exhibiting defining stem cell characteristics of multi-potency and self-renewal. These factors were shown to act co-operatively as expression of each factor individually was

\*Correspondence: SmalleyMJ@Cardiff.ac.uk

<sup>2</sup>European Cancer Stem Cell Institute, Cardiff School of Biosciences, Cardiff University Life Sciences Building, Museum Avenue, Cardiff CF10 3AX, UK  
Full list of author information is available at the end of the article

not sufficient to confer stem cell traits on fully differentiated cells.

Ectopic *Sox9* expression alone in differentiated cells produced cells with luminal progenitor characteristics, being able to proliferate *in vitro* and form hollow acini in three-dimensional culture conditions. Ectopic expression of *Slug* was able to convert luminal progenitor cells, which endogenously express *Sox9*, into cells with stem cell activity. In contrast, stem cell activity was enhanced in basal cells, which endogenously express *Slug*, by forced expression of *Sox9*. Interestingly, when *Sox9* was ectopically expressed in basal cells at the same time as *Slug* was knocked down, the cells acquired a luminal progenitor-like phenotype *in vitro*. Gene expression analysis of differentiated cells ectopically expressing *Slug* or *Sox9* showed upregulation of basal or luminal progenitor-associated genes, respectively, and both signatures were upregulated upon *Slug/Sox9* co-expression. The authors postulated that *Slug* and *Sox9* regulate basal and luminal lineage programmes, respectively. Each confers distinct biological properties on the cell, but both are required for MaSC function.

Given the important potential link between MaSCs and breast cancer stem cells, the study next looked at the effects of *Slug* and *Sox9* on tumour-initiating potential and metastasis. The human breast cancer cell line MDA-MB-231 expresses both *Slug* and *Sox9*, forms tumours upon subcutaneous injection and metastasises to the lung upon tail vein injection into NOD/SCID mice. Knockdown of *Sox9* reduced the tumour-initiating potential of the cells by over 70-fold following subcutaneous injection. In contrast, knockdown of *Slug* did not affect tumour initiation but the resulting tumours were sixfold smaller than those in controls. In a metastasis assay, *Slug* or *Sox9* knockdown reduced lung metastases by fivefold and 40-fold, respectively. In non-metastatic MCF7ras human breast cancer cells implanted orthotopically into NOD/SCID mice, there was a significant increase in macrometastases from primary tumours ectopically co-expressing *Slug* and *Sox9*. Finally, in a tissue microarray of 306 clinical breast cancer samples, patients with primary tumours expressing high levels of both *Slug* and *Sox9* had a significantly lower overall survival rate. The authors conclude that human breast cancer stem cells are controlled by the same master regulators governing the murine MaSC state.

### Stem cell enhancement or altered cell fate?

Many studies have now demonstrated that transplantation potential in the cleared mammary fat pad assay is most highly enriched in the basal cell layer of the mammary epithelium [2-4]. If this transplantation potential can be serially propagated, we call this mammary epithelial stem cell potential. However, both the potential plasticity of

the normal mammary epithelium and the inductive potential of the mammary environment are increasingly well recognised.

The work of Guo and colleagues is a very important step forward in understanding the molecular regulation of this plasticity and how it can promote mammary stem cell potential, but it will be difficult to disentangle these phenomena. *Slug* promotes the basal phenotype when ectopically expressed in luminal progenitors. *Sox9* promotes luminal progenitor activity in differentiated luminal cells and stem cell activity in differentiated basal cells. Together, *Slug* and *Sox9* determine stem cell-like activity in the mammary epithelium. Whether the basal phenotype is required for this activity (perhaps by determining response to the microenvironment) or whether changes in underlying self-renewal processes are independent of phenotypic changes remains unclear. Perhaps both scenarios are occurring, with *Sox9* priming cellular self-renewal to respond to niche interactions that are made possible by the transcriptional programme driven by *Slug* activity.

Finally, as Guo and colleagues note themselves, in the mammary epithelium 'basal' does not necessarily mean 'stem cell'; and indeed the majority of basal mammary cells are not stem cells but rather are myoepithelial cells. Myoepithelial cells are, in terms of both function and identity, a cross between epithelial cells and mesenchymal cells and they express numerous EMT markers as part of their normal biology [10]. It is therefore wrong to characterise a luminal to basal phenotypic switch as EMT. Rather, expression of 'mesenchymal markers' is part of the normal differentiative programme of the mammary epithelium, and this paper must be understood in that context.

#### Abbreviations

EMT, epithelial-mesenchymal transition; MaSC, mammary stem cell; NOD/SCID, nonobese diabetic/severe combined immunodeficiency.

#### Competing interests

The authors declare that they have no competing interests.

#### Acknowledgements

KS is supported by an ICR/MRC Studentship. MJS is supported by the European Cancer Stem Cell Research Institute and Cardiff University.

#### Author details

<sup>1</sup>Breakthrough Breast Cancer Research Centre, The Institute of Cancer Research, 237 Fulham Road, London SW3 6JB, UK. <sup>2</sup>European Cancer Stem Cell Research Institute, Cardiff School of Biosciences, Cardiff University Life Sciences Building, Museum Avenue, Cardiff CF10 3AX, UK.

Published: 10 September 2012

#### References

1. Deome KB, Faulkin LJ, Jr, Bern HA, Blair PB: Development of mammary tumors from hyperplastic alveolar nodules transplanted into gland-free mammary fat pads of female C3H mice. *Cancer Res* 1959, **19**:515-520.
2. Stingl J, Eirew P, Ricketson I, Shackleton M, Vaillant F, Choi D, Li HI, Eaves CJ: Purification and unique properties of mammary epithelial stem cells. *Nature* 2006, **439**:993-997.

3. Shackleton M, Vaillant F, Simpson KJ, Stingl J, Smyth GK, Asselin-Labat ML, Wu L, Lindeman GJ, Visvader JE: **Generation of a functional mammary gland from a single stem cell.** *Nature* 2006, **439**:84-88.
4. Sleeman KE, Kendrick H, Ashworth A, Isacke CM, Smalley MJ: **CD24 staining of mouse mammary gland cells defines luminal epithelial, myoepithelial/basal and non-epithelial cells.** *Breast Cancer Res* 2006, **8**:R7.
5. Regan JL, Kendrick H, Magnay FA, Vafaizadeh V, Groner B, Smalley MJ: **c-Kit is required for growth and survival of the cells of origin of Brca1-mutation-associated breast cancer.** *Oncogene* 2012, **31**:869-883.
6. Mani SA, Guo W, Liao MJ, Eaton EN, Ayyanan A, Zhou AY, Brooks M, Reinhard F, Zhang CC, Shipitsin M, Campbell LL, Polyak K, Brisken C, Yang J, Weinberg RA: **The epithelial–mesenchymal transition generates cells with properties of stem cells.** *Cell* 2008, **133**:704-715.
7. Morel AP, Lièvre M, Thomas C, Hinkal G, Ansieau S, Puisieux A: **Generation of breast cancer stem cells through epithelial–mesenchymal transition.** *Plos One* 2008, **3**:e2888.
8. Thiery JP, Acloque H, Huang RYJ, Nieto MA: **Epithelial–mesenchymal transitions in development and disease.** *Cell* 2009, **139**:871-890.
9. Guo W, Keckesova Z, Donaher JL, Shibue T, Tischler V, Reinhardt F, Itzkovitz S, Noske A, Zürcher-Härdi U, Bell G, Tam WL, Mani SA, van Oudenaarden A, Weinberg RA: **Slug and Sox9 cooperatively determine the mammary stem cell state.** *Cell* 2012, **148**:1015-1028.
10. Kendrick H, Regan JL, Magnay FA, Grigoriadis A, Mitsopoulos C, Zvelebil M, Smalley MJ: **Transcriptome analysis of mammary epithelial subpopulations identifies novel determinants of lineage commitment and cell fate.** *BMC Genomics* 2008, **9**:591.

doi:10.1186/bcr3188

**Cite this article as:** Soady K, Smalley MJ: **Slugging their way to immortality: driving mammary epithelial cells into a stem cell-like state.** *Breast Cancer Research* 2012, **14**:319.